arXiv:1208.4584v1 [math.AG] 22 Aug 2012

MONODROMIES AT INFINITY OF NON-TAME POLYNOMIALS

KIYOSHI TAKEUCHI AND MIHAI TIBĂR

Abstract. We study monodromies at infinity of non-tame polynomials. In particular we prove some formulae which express their Jordan normal forms in terms of the Newton polyhedra at infinity.

1. Introduction

For a polynomial map $f\colon \mathbb{C}^n \longrightarrow \mathbb{C}$, it is well-known that there exists a finite subset $B \subset \mathbb{C}$ such that the restriction

$$\mathbb{C}^n \setminus f^{-1}(B) \longrightarrow \mathbb{C} \setminus B \tag{1.1}$$

of f is a locally trivial fibration. We denote by B_f the smallest subset $B \subset \mathbb{C}$ satisfying this condition. We call elements of B_f bifurcation points of f. Let $C_R = \{x \in \mathbb{C} \mid |x| = R\}$ $(R \gg 0)$ be a sufficiently large circle in $\mathbb{C}$ such that $B_f \subset \{x \in \mathbb{C} \mid |x| < R\}$. Then by restricting the locally trivial fibration $\mathbb{C}^n \setminus f^{-1}(B_f) \longrightarrow \mathbb{C} \setminus B_f$ to C_R we obtain a geometric monodromy automorphism $\Phi_f^\infty\colon f^{-1}(R) \xrightarrow{\sim} f^{-1}(R)$ and the linear maps

$$\Phi_j^\infty\colon H^j(f^{-1}(R);\mathbb{C}) \xrightarrow{\sim} H^j(f^{-1}(R);\mathbb{C}) \quad (j = 0, 1, \dots) \tag{1.2}$$

associated to it, where the orientation of C_R is taken to be counter-clockwise as usual. We call Φ_j^∞'s the (cohomological) monodromies at infinity of f. In the last few decades many mathematicians studied Φ_j^∞'s from various points of view. In particular in [1] Broughton proved that if f is tame at infinity we have the concentration

$$H^j(f^{-1}(R);\mathbb{C}) = 0 \quad (j \neq 0, n-1) \tag{1.3}$$

for the generic fiber $f^{-1}(R)$ $(R \gg 0)$ of f. This implies that in the tame case Φ_{n-1}^∞ is the only non-trivial monodromy at infinity to study. In this case Libgober-Sperber [14] obtained a beautiful formula which expresses the semisimple part (i.e. the eigenvalues) of Φ_{n-1}^∞ in terms of the Newton polyhedron at infinity of f (see [16] for its generalizations). Recently in [18] (see also [7]) we proved also formulae for its nilpotent part (i.e. its Jordan normal form) by introducing the motivic Milnor fiber at infinity of f. However if we do not assume the tameness at infinity of f, Broughton's theorem does not hold in general. In the non-tame case, even if we try to refine its generalization in [28] we can not prove the above cohomological concentration on the generic fiber of f. Indeed we can easily find a lot of counterexamples. This leads us to face an essential difficulty in applying our methods in [18] and [7] to non-tame polynomials.

In this paper, we partially overcome this problem by showing that the desired cohomological concentration holds for the generalized eigenspaces of Φ_j^∞ for "good" eigenvalues.

2000 *Mathematics Subject Classification.* 14E18, 14M25, 32C38, 32S35, 32S40.

Namely if we avoid some "bad" eigenvalues associated to f, we can successfully generalize the results in [18] to non-tame polynomials and completely determine the Jordan normal forms of Φ_{n-1}^{∞}. Note that in the study of (local) Milnor monodromies also Dimca-Saito [6] avoid some bad eigenvalues to deal with non-isolated singular points. In this sense, some of our results in this paper can be considered as the global analogues of those in [6]. In order to explain our results more precisely, let us recall some basic definitions. For $f(x) = \sum_{v \in \mathbb{Z}_+^n} a_v x^v$ $(a_v \in \mathbb{C})$ we call the convex hull of $\mathrm{supp} f = \{v \in \mathbb{R}_+^n \mid a_v \neq 0\}$ in $\mathbb{R}^n$ the Newton polytope of f and denote it by $NP(f)$. The convex hull $\Gamma_\infty(f) \subset \mathbb{R}_+^n$ of $\{0\} \cup NP(f)$ in $\mathbb{R}^n$ is called the Newton polyhedron at infinity of f. Assume that $\dim \Gamma_\infty(f) = n$ and f is non-degenerate at infinity (see Section 2 for the definition).

Definition 1.1. *We say the f is convenient if $\Gamma_\infty(f)$ intersects the positive part of the i-th axis of $\mathbb{R}^n$ for any $1 \leq i \leq n$.*

If f is convenient and non-degenerate at infinity, then by a result of Broughton [1] it is tame at infinity. However here we do not assume that f is convenient. Let us assume that f is not convenient. In Section 2 by using the dual fan of $\Gamma_\infty(f)$ we define a finite subset $A_f \subset \mathbb{C}$ of "bad" eigenvalues of Φ_j^{∞}'s which we call atypical engenvalues of f. Then the following result plays a key role in this paper. For $\lambda \in \mathbb{C}$ and $j \in \mathbb{Z}$ let $H^j(f^{-1}(R); \mathbb{C})_\lambda \subset H^j(f^{-1}(R); \mathbb{C})$ be the generalized eigenspace for the eigenvalue λ of the monodromy at infinity $\Phi_j^{\infty} \colon H^j(f^{-1}(R); \mathbb{C}) \xrightarrow{\sim} H^j(f^{-1}(R); \mathbb{C})$ $(R \gg 0)$.

Theorem 1.2. *In the situation as above, for any non-atypical eigenvalue $\lambda \notin A_f$ of f we have the concentration*

$$H^j(f^{-1}(R); \mathbb{C})_\lambda \simeq 0 \qquad (j \neq n-1) \tag{1.4}$$

for the generic fiber $f^{-1}(R) \subset \mathbb{C}^n$ $(R \gg 0)$ of f.

We prove this theorem by refining the proof of Sabbah's theorem [27, Theorem 13.1] in our situation. More precisely we construct a nice compactification $\widetilde{X_\Sigma}$ of $\mathbb{C}^n$ and study the "horizontal" divisors at infinity for f in $\widetilde{X_\Sigma} \setminus \mathbb{C}^n$ carefully to prove the concentration. With Theorem 1.2 at hand, by using the results in [18, Section 2] we can easily prove the analogues of [18, Theorems 5.9, 5.14 and 5.16] for the non-tame polynomial f and completely determine the λ-part of the Jordan normal form of Φ_{n-1}^{∞} for any $\lambda \notin A_f$. Here let us introduce only that of [18, Theorem 5.9]. Denote by $\mathrm{Cone}_\infty(f)$ the closed cone $\mathbb{R}_+\Gamma_\infty(f) \subset \mathbb{R}_+^n$ generated by $\Gamma_\infty(f)$. Let $q_1, \dots, q_l$ (resp. $\gamma_1, \dots, \gamma_{l'}$) be the 0-dimensional (resp. 1-dimensional) faces of $\Gamma_\infty(f)$ such that $q_i \in \mathrm{Int}(\mathrm{Cone}_\infty(f))$ (resp. the relative interior $\mathrm{rel.int}(\gamma_i)$ of γ_i is contained in $\mathrm{Int}(\mathrm{Cone}_\infty(f))$). For each q_i (resp. γ_i), denote by $d_i > 0$ (resp. $e_i > 0$) the lattice distance $\mathrm{dist}(q_i, 0)$ (resp. $\mathrm{dist}(\gamma_i, 0)$) of it from the origin $0 \in \mathbb{R}^n$. For $1 \leq i \leq l'$, let Δ_i be the convex hull of $\{0\} \sqcup \gamma_i$ in $\mathbb{R}^n$. Then for $\lambda \neq 1$ and $1 \leq i \leq l'$ such that $\lambda^{e_i} = 1$ we set

$$n(\lambda)_i = \sharp\{v \in \mathbb{Z}^n \cap \mathrm{rel.int}(\Delta_i) \mid \mathrm{ht}(v, \gamma_i) = k\} + \sharp\{v \in \mathbb{Z}^n \cap \mathrm{rel.int}(\Delta_i) \mid \mathrm{ht}(v, \gamma_i) = e_i - k\}, \tag{1.5}$$

where k is the minimal positive integer satisfying $\lambda = \zeta_{e_i}^k$ (we set $\zeta_d := \exp(2\pi\sqrt{-1}/d) \in \mathbb{C}$) and for $v \in \mathbb{Z}^n \cap \mathrm{rel.int}(\Delta_i)$ we denote by $\mathrm{ht}(v, \gamma_i)$ the lattice height of v from the base γ_i of Δ_i. Then we have the following analogue of [18, Theorem 5.9].

Theorem 1.3. *In the situation as above, for any* $\lambda \notin A_f$ *we have*

1. *The number of the Jordan blocks for the eigenvalue* λ *with the maximal possible size* n *in* $\Phi_{n-1}^{\infty}\colon H^{n-1}(f^{-1}(R);\mathbb{C}) \xrightarrow{\sim} H^{n-1}(f^{-1}(R);\mathbb{C})$ $(R \gg 0)$ *is equal to* $\sharp\{q_i \mid \lambda^{d_i} = 1\}$.
2. *The number of the Jordan blocks for the eigenvalue* λ *with the second maximal possible size* $n-1$ *in* Φ_{n-1}^{∞} *is equal to* $\sum_{i\colon \lambda^{e_i}=1} n(\lambda)_i$.

Similarly we can treat also the following important monodromies associated to f. Let $b \in B_f$ be a bifurcation point of f. Choose sufficiently small $\varepsilon > 0$ such that

$$B_f \cap \{x \in \mathbb{C} \mid |x-b| \leq \varepsilon\} = \{b\} \tag{1.6}$$

and set $C_\varepsilon(b) = \{x \in \mathbb{C} \mid |x-b| = \varepsilon\} \subset \mathbb{C}$. Then we obtain a locally trivial fibration $f^{-1}(C_\varepsilon(b)) \longrightarrow C_\varepsilon(b)$ over the small circle $C_\varepsilon(b) \subset \mathbb{C}$ and the monodromy automorphisms

$$\Phi_j^b\colon H^j(f^{-1}(b+\varepsilon);\mathbb{C}) \xrightarrow{\sim} H^j(f^{-1}(b+\varepsilon);\mathbb{C}) \quad (j = 0, 1, \ldots) \tag{1.7}$$

around the central fiber $f^{-1}(b) \subset \mathbb{C}^n$ associated to it. In Section 6 we apply our methods to the Jordan normal forms of Φ_j^b's. Since we assume here that f is not convenient, for some $b \in B_f$ the central fiber $f^{-1}(b) \subset \mathbb{C}^n$ may have "singularities at infinity" studied by Némethi-Zaharia [22] and Zaharia [35] etc. Even in such cases, we can define a finite subset $A_{f,b}^{\circ} \subset \mathbb{C}$ of "bad" eigenvalues of Φ_j^b's and completely determine the λ-part of the Jordan normal form of Φ_{n-1}^b for any $\lambda \notin A_{f,b}^{\circ}$. In fact we obtain these results more generally for polynomial maps $f : U \longrightarrow \mathbb{C}$ of affine algebraic varieties U. See Section 6 for the details.

2. MONODROMIES AT INFINITY

Let $f(x)$ be a polynomial on $\mathbb{C}^n$. Then as we explained in Introduction, there exist a locally trivial fibration $\mathbb{C}^n \setminus f^{-1}(B_f) \longrightarrow \mathbb{C} \setminus B_f$ and the linear maps

$$\Phi_j^{\infty}\colon H^j(f^{-1}(R);\mathbb{C}) \xrightarrow{\sim} H^j(f^{-1}(R);\mathbb{C}) \quad (j = 0, 1, \ldots) \tag{2.1}$$

$(R \gg 0)$ associated to it. To study the monodromies at infinity Φ_j^{∞}, we often impose the following natural condition.

Definition 2.1 ([13]). *Let* $\partial f\colon \mathbb{C}^n \longrightarrow \mathbb{C}^n$ *be the map defined by* $\partial f(x) = (\partial_1 f(x), \ldots, \partial_n f(x))$. *Then we say that* f *is tame at infinity if the restriction* $(\partial f)^{-1}(B(0;\varepsilon)) \longrightarrow B(0;\varepsilon)$ *of* ∂f *to a sufficiently small ball* $B(0;\varepsilon)$ *centered at the origin* $0 \in \mathbb{C}^n$ *is proper.*

The following result is fundamental in the study of monodromies at infinity.

Theorem 2.2 (Broughton [1] and Siersma-Tibăr [28]). *Assume that* f *is tame at infinity (more generally, with isolated* $\mathcal{W}$*-singularities, see* [28]*). Then the generic fiber* $f^{-1}(c)$ $(c \in \mathbb{C} \setminus B_f)$ *has the homotopy type of the bouquet of* $(n-1)$*-spheres. In particular, we have*

$$H^j(f^{-1}(c);\mathbb{C}) = 0 \quad (j \neq 0, n-1). \tag{2.2}$$

□

By this theorem if f is tame at infinity, then Φ_{n-1}^{∞} is the only non-trivial monodromy at infinity.

Definition 2.3 ([14]). *We call the convex hull of* $\{0\} \cup NP(f)$ *in* $\mathbb{R}^n$ *the Newton polyhedron at infinity of* f *and denote it by* $\Gamma_\infty(f)$.

Definition 2.4 ([13]). *We say that the polynomial* $f(x) = \sum_{v \in \mathbb{Z}_+^n} a_v x^v$ $(a_v \in \mathbb{C})$ *is non-degenerate at infinity if for any face* γ *of* $\Gamma_\infty(f)$ *such that* $0 \notin \gamma$ *the complex hypersurface* $\{x \in (\mathbb{C}^*)^n \mid f_\gamma(x) = 0\}$ *in* $(\mathbb{C}^*)^n$ *is smooth and reduced, where we set* $f_\gamma(x) = \sum_{v \in \gamma \cap \mathbb{Z}_+^n} a_v x^v$.

If f is convenient and non-degenerate at infinity, then by a result of Broughton [1] it is tame at infinity. However in this paper, we do not assume that f is convenient. If $\dim \Gamma_\infty(f) = n$, to the n-dimensional integral polytope $\Gamma_\infty(f)$ in $\mathbb{R}^n$ we can naturally associate a subdivision of (the dual vector space of) $\mathbb{R}^n$ into rational convex cones as follows. For an element $u \in \mathbb{R}^n$ of (the dual vector space of) $\mathbb{R}^n$ define the supporting face $\gamma_u \prec \Gamma_\infty(f)$ of u in $\Gamma_\infty(f)$ by

$$\gamma_u = \left\{ v \in \Gamma_\infty(f) \mid \langle u, v \rangle = \min_{w \in \Gamma_\infty(f)} \langle u, w \rangle \right\}. \tag{2.3}$$

Then we introduce an equivalence relation $\sim$ on (the dual vector space of) $\mathbb{R}^n$ by $u \sim u' \iff \gamma_u = \gamma_{u'}$. We can easily see that for any face $\gamma \prec \Gamma_\infty(f)$ of $\Gamma_\infty(f)$ the closure of the equivalence class associated to γ in $\mathbb{R}^n$ is an $(n - \dim \gamma)$-dimensional rational convex polyhedral cone $\sigma(\gamma)$ in $\mathbb{R}^n$. Moreover the family $\{\sigma(\gamma) \mid \gamma \prec \Gamma_\infty(f)\}$ of cones in $\mathbb{R}^n$ thus obtained is a subdivision of $\mathbb{R}^n$ and satisfies the axiom of fans (see [8] and [23] etc.). We call it the dual fan of $\Gamma_\infty(f)$.

Definition 2.5. *Assume that* $\dim \Gamma_\infty(f) = n$. *Then we say that a face* $\gamma \prec \Gamma_\infty(f)$ *is atypical if* $0 \in \gamma$ *and the cone* $\sigma(\gamma)$ *which corresponds to it in the dual fan of* $\Gamma_\infty(f)$ *is not contained in the first quadrant* $\mathbb{R}_+^n$ *in* $\mathbb{R}^n$.

Example 2.6. *Let* $n = 3$ *and consider a non-convenient polynomial* $f(x, y, z)$ *on* $\mathbb{C}^3$ *whose Newton polyhedron at infinity* $\Gamma_\infty(f)$ *is the convex hull of the points* $(2, 0, 0), (2, 2, 0), (2, 2, 3) \in \mathbb{R}_+^3$ *and the origin* $0 = (0, 0, 0) \in \mathbb{R}^3$. *Then the line segment whose vertices are the point* $(2, 2, 0)$ *and the origin* $0 \in \mathbb{R}^3$ *is an atypical face of* $\Gamma_\infty(f)$. *However the triangle whose vertices are the points* $(2, 0, 0)$, $(2, 2, 0)$ *and the origin* $0 \in \mathbb{R}^3$ *is not so.*

Example 2.7. *Let* $n = 3$ *and consider a non-convenient polynomial* $f(x, y, z)$ *on* $\mathbb{C}^3$ *whose Newton polyhedron at infinity* $\Gamma_\infty(f)$ *is the convex hull of the points* $(2, 0, 0), (0, 2, 0), (1, 1, 2) \in \mathbb{R}_+^3$ *and the origin* $0 = (0, 0, 0) \in \mathbb{R}^3$. *Then the line segment connecting the point* $(2, 0, 0)$ *and the origin* $0 \in \mathbb{R}^3$ *is an atypical face of* $\Gamma_\infty(f)$.

For a subset $S \subset \{1, 2, \dots, n\}$ we define a coordinate subspace $\mathbb{R}^S \simeq \mathbb{R}^{|S|}$ of $\mathbb{R}^n$ by

$$\mathbb{R}^S = \{v = (v_1, \dots, v_n) \in \mathbb{R}^n \mid v_i = 0 \quad \text{for any} \quad i \notin S\}. \tag{2.4}$$

The following lemma should be obvious.

Lemma 2.8. *Assume that* $\dim \Gamma_\infty(f) = n$ *and let* $\gamma \prec \Gamma_\infty(f)$ *be a face of* $\Gamma_\infty(f)$ *such that* $0 \in \gamma$. *Let* $\mathbb{R}^S \subset \mathbb{R}^n$ *be the minimal coordinate subspace of* $\mathbb{R}^n$ *containing* γ *and assume that* $\dim \gamma < \dim \mathbb{R}^S = |S|$. *Then* γ *is an atypical face of* $\Gamma_\infty(f)$. □

By this lemma we can easily prove the following proposition.

Proposition 2.9. *Assume that* $\dim \Gamma_\infty(f) = n$ *and a face* $\gamma \prec \Gamma_\infty(f)$ *of* $\Gamma_\infty(f)$ *such that* $0 \in \gamma$ *is not atypical. Let* $\mathbb{R}^S \subset \mathbb{R}^n$ *be the minimal coordinate subspace of* $\mathbb{R}^n$ *containing* γ. *Then we have* $\dim \gamma = \dim \mathbb{R}^S = |S|$ *and there exist exactly* $n - |S|$ *facets i.e.* $(n-1)$-*dimensional faces* $\Gamma_i \prec \Gamma_\infty(f)$ $(i \notin S)$ *of* $\Gamma_\infty(f)$ *containing* γ. *Moreover they are explicitly given by*

$$\Gamma_i = \Gamma_\infty(f) \cap \{v = (v_1, \dots, v_n) \in \mathbb{R}^n \mid v_i = 0\} \qquad (i \notin S). \tag{2.5}$$

□

Definition 2.10. *We say that a face* $\gamma \prec \Gamma_\infty(f)$ *of* $\Gamma_\infty(f)$ *is at infinity if* $0 \notin \gamma$.

Definition 2.11. *Let* $\gamma \prec \Gamma_\infty(f)$ *be a face at infinity of* $\Gamma_\infty(f)$. *Then we say that* γ *is admissible if it is not contained in any atypical face of* $\Gamma_\infty(f)$.

For a face at infinity $\gamma \prec \Gamma_\infty(f)$ of $\Gamma_\infty(f)$, let Δ_γ be the convex hull of $\{0\} \sqcup \gamma$ in $\mathbb{R}^n$. Denote by $\mathbb{L}(\Delta_\gamma)$ the $(\dim \gamma + 1)$-dimensional linear subspace of $\mathbb{R}^n$ spanned by Δ_γ and consider the lattice $M_\gamma = \mathbb{Z}^n \cap \mathbb{L}(\Delta_\gamma) \simeq \mathbb{Z}^{\dim \gamma + 1}$ in it. Then there exists a unique non-zero primitive vector u_γ in its dual lattice which takes its maximum in Δ_γ exactly on $\gamma \prec \Delta_\gamma$:

$$\gamma = \left\{ v \in \Delta_\gamma \mid \langle u_\gamma, v \rangle = \max_{w \in \Delta_\gamma} \langle u_\gamma, w \rangle \right\}. \tag{2.6}$$

We set

$$d_\gamma = \max_{w \in \Delta_\gamma} \langle u_\gamma, w \rangle \in \mathbb{Z}_{>0} \tag{2.7}$$

and call it the lattice distance of γ from the origin $0 \in \mathbb{R}^n$. The following definition will be used in Section 4.

Definition 2.12. *Assume that* $\dim \Gamma_\infty(f) = n$. *Then we say that a complex number* $\lambda \in \mathbb{C}$ *is an atypical eigenvalue of* f *if there exists a non-admissible face at infinity* $\gamma \prec \Gamma_\infty(f)$ *of* $\Gamma_\infty(f)$ *such that* $\lambda^{d_\gamma} = 1$. *We denote by* $A_f \subset \mathbb{C}$ *the set of the atypical eigenvalues of* f.

Let $\overrightarrow{e_i} = {}^t(0, \dots, 0, \overset{i}{1}, 0, \dots, 0) \in \mathbb{R}^n$ $(i = 1, 2, \dots, n)$ be the standard basis of $\mathbb{R}^n$. The following result will be used in Section 3.

Proposition 2.13. *Assume that* $\dim \Gamma_\infty(f) = n$ *and let* $\gamma \prec \Gamma_\infty(f)$ *be a face at infinity of* $\Gamma_\infty(f)$. *Let* $\mathbb{R}^S \subset \mathbb{R}^n$ *be the minimal coordinate subspace of* $\mathbb{R}^n$ *containing* γ *and* $\sigma \subset \mathbb{R}^n$ *the cone in the dual fan of* $\Gamma_\infty(f)$ *which corresponds to* $\gamma \prec \Gamma_\infty(f)$. *Then* γ *is admissible if and only if there exist some integral vectors* $\overrightarrow{f_1}, \dots, \overrightarrow{f_k} \in (\mathbb{R}^n \setminus \mathbb{R}^n_+) \cap \mathbb{Z}^n$ *such that*

$$\min_{v \in \Gamma_\infty(f)} \langle \overrightarrow{f_j}, v \rangle < 0 \qquad (1 \leq j \leq k) \tag{2.8}$$

and

$$\sigma=(\sum_{i\notin S}\mathbb{R}_+\overrightarrow{e_i})+(\sum_{j=1}^{k}\mathbb{R}_+\overrightarrow{f_j}). \tag{2.9}$$

□

3. Motivic Milnor fibers at infinity

From now, following Denef-Loeser [3] and [4] we introduce motivic reincarnations of global (Milnor) fibers of polynomial maps f. For the details, see Matsui-Takeuchi [18], Esterov-Takeuchi [7] and Raibaut [25]. First, take a smooth compactification X of $\mathbb{C}^n$. Then by eliminating the points of indeterminacy of the meromorphic extension of f to X we obtain a commutative diagram

$$\begin{tikzcd} \mathbb{C}^n \arrow[r, "\iota"] \arrow[d, "f"'] & \widetilde{X} \arrow[d, "g"] \\ \mathbb{C} \arrow[r, "j"'] & \mathbb{P}^1 \end{tikzcd} \tag{3.1}$$

such that horizontal arrows are open embeddings, g is a proper holomorphic map and $\widetilde{X}\setminus\mathbb{C}^n$, $Y:=g^{-1}(\infty)$ are normal crossing divisors in $\widetilde{X}$. Take a local coordinate h of $\mathbb{P}^1$ in a neighborhood of $\infty\in\mathbb{P}^1$ such that $\infty=\{h=0\}$ and set $\widetilde{g}=h\circ g$. Note that $\widetilde{g}$ is a holomorphic function defined on a neighborhood of the closed subvariety $Y=\widetilde{g}^{-1}(0)=g^{-1}(\infty)\subset\widetilde{X}\setminus\mathbb{C}^n$ of $\widetilde{X}$. Then for $R\gg 0$ we have

$$H^j_c(f^{-1}(R);\mathbb{C})\simeq H^j(Y;\psi_{\widetilde{g}}(\iota_!\mathbb{C}_{\mathbb{C}^n})). \tag{3.2}$$

Let us define an open subset Ω of $\widetilde{X}$ by

$$\Omega=\mathrm{Int}(\iota(\mathbb{C}^n)\sqcup Y) \tag{3.3}$$

and set $U=\Omega\cap Y$. Then U (resp. the complement of Ω in $\widetilde{X}$) is a normal crossing divisor in Ω (resp. $\widetilde{X}$). Hence we can easily prove the isomorphisms

$$H^j(Y;\psi_{\widetilde{g}}(\iota_!\mathbb{C}_{\mathbb{C}^n}))\simeq H^j(Y;\psi_{\widetilde{g}}(\iota'_!\mathbb{C}_{\Omega}))\simeq H^j_c(U;\psi_{\widetilde{g}}(\mathbb{C}_{\widetilde{X}})), \tag{3.4}$$

where $\iota'\colon\Omega\hookrightarrow\widetilde{X}$ is the inclusion. Now let $E_1,E_2,\dots,E_k$ be the irreducible components of the normal crossing divisor $U=\Omega\cap Y$ in $\Omega\subset\widetilde{X}$. For each $1\le i\le k$, let $b_i>0$ be the order of the zero of $\widetilde{g}$ along E_i. For a non-empty subset $I\subset\{1,2,\dots,k\}$, let us set

$$E_I=\bigcap_{i\in I}E_i,\qquad E_I^\circ=E_I\setminus\bigcup_{i\notin I}E_i \tag{3.5}$$

and $d_I=\gcd(b_i)_{i\in I}>0$. Then, as in [4, Section 3.3], we can construct an unramified Galois covering $\widetilde{E_I^\circ}\longrightarrow E_I^\circ$ of E_I° as follows. First, for a point $p\in E_I^\circ$ we take an affine open neighborhood $W\subset\Omega\setminus(\cup_{i\notin I}E_i)$ of p on which there exist regular functions ξ_i $(i\in I)$ such that $E_i\cap W=\{\xi_i=0\}$ for any $i\in I$. Then on W we have $\widetilde{g}=\widetilde{g_{1,W}}(\widetilde{g_{2,W}})^{d_I}$, where we set $\widetilde{g_{1,W}}=\widetilde{g}\prod_{i\in I}\xi_i^{-b_i}$ and $\widetilde{g_{2,W}}=\prod_{i\in I}\xi_i^{\frac{b_i}{d_I}}$. Note that $\widetilde{g_{1,W}}$ is a unit on W and

$\widetilde{g_{2,W}}\colon W \longrightarrow \mathbb{C}$ is a regular function. It is easy to see that E_I° is covered by such affine open subsets W of $\Omega \setminus (\cup_{i\notin I} E_i)$. Then as in [4, Section 3.3] by gluing the varieties

$$\widetilde{E_{I,W}^\circ} = \{(t,z) \in \mathbb{C}^* \times (E_I^\circ \cap W) \mid t^{d_I} = (\widetilde{g_{1,W}})^{-1}(z)\} \tag{3.6}$$

together in the following way, we obtain the variety $\widetilde{E_I^\circ}$ over E_I°. If W' is another such open subset and $\widetilde{g} = \widetilde{g_{1,W'}}(\widetilde{g_{2,W'}})^{d_I}$ is the decomposition of $\widetilde{g}$ on it, we patch $\widetilde{E_{I,W}^\circ}$ and $\widetilde{E_{I,W'}^\circ}$ by the morphism $(t,z) \longmapsto (\widetilde{g_{2,W'}}(z)(\widetilde{g_{2,W}})^{-1}(z)\cdot t, z)$ defined over $W \cap W'$. Now for $d \in \mathbb{Z}_{>0}$, let $\mu_d \simeq \mathbb{Z}/\mathbb{Z}d$ be the multiplicative group consisting of the d-roots in $\mathbb{C}$. We denote by $\hat{\mu}$ the projective limit $\varprojlim_d \mu_d$ of the projective system $\{\mu_i\}_{i\geq 1}$ with morphisms $\mu_{id} \longrightarrow \mu_i$ given by $t \longmapsto t^d$. Then the unramified Galois covering $\widetilde{E_I^\circ}$ of E_I° admits a natural μ_{d_I}-action defined by assigning the automorphism $(t,z) \longmapsto (\zeta_{d_I} t, z)$ of $\widetilde{E_I^\circ}$ to the generator $\zeta_{d_I} := \exp(2\pi\sqrt{-1}/d_I) \in \mu_{d_I}$. Namely the variety $\widetilde{E_I^\circ}$ is equipped with a good $\hat{\mu}$-action in the sense of [4, Section 2.4]. Following the notations in [4], denote by $\mathcal{M}_{\mathbb{C}}^{\hat{\mu}}$ the ring obtained from the Grothendieck ring $\mathrm{K}_0^{\hat{\mu}}(\mathrm{Var}_{\mathbb{C}})$ of varieties over $\mathbb{C}$ with good $\hat{\mu}$-actions by inverting the Lefschetz motive $\mathbb{L} \simeq \mathbb{C} \in \mathrm{K}_0^{\hat{\mu}}(\mathrm{Var}_{\mathbb{C}})$. Recall that $\mathbb{L} \in \mathrm{K}_0^{\hat{\mu}}(\mathrm{Var}_{\mathbb{C}})$ is endowed with the trivial action of $\hat{\mu}$.

Definition 3.1 ([18] and [25]). *We define the motivic Milnor fiber at infinity $\mathcal{S}_f^\infty$ of the polynomial map $f\colon \mathbb{C}^n \longrightarrow \mathbb{C}$ by*

$$\mathcal{S}_f^\infty = \sum_{I\neq\emptyset} (1-\mathbb{L})^{|I|-1}[\widetilde{E_I^\circ}] \in \mathcal{M}_{\mathbb{C}}^{\hat{\mu}}. \tag{3.7}$$

Remark 3.2. *By Guibert-Loeser-Merle* [10, Theorem 3.9]*, the motivic Milnor fiber at infinity $\mathcal{S}_f^\infty$ of f does not depend on the compactification X of $\mathbb{C}^n$. This fact was informed to us by Schürmann (a private communication) and Raibaut* [25].

As in [4, Section 3.1.2 and 3.1.3], we denote by $\mathrm{HS}^{\mathrm{mon}}$ the abelian category of Hodge structures with a quasi-unipotent endomorphism. Then, to the object $\psi_h(j_! Rf_! \mathbb{C}_{\mathbb{C}^n}) \in \mathbf{D}_c^b(\{\infty\})$ and the semisimple part of the monodromy automorphism acting on it, we can associate an element

$$[H_f^\infty] \in \mathrm{K}_0(\mathrm{HS}^{\mathrm{mon}}) \tag{3.8}$$

as in [3] and [4]. Recall that the weight filtrations of $H^j\psi_h(j_! Rf_! \mathbb{C}_{\mathbb{C}^n})$ in the construction of $[H_f^\infty]$ are "relative" monodromy filtrations. To describe the element $[H_f^\infty] \in \mathrm{K}_0(\mathrm{HS}^{\mathrm{mon}})$ in terms of $\mathcal{S}_f^\infty \in \mathcal{M}_{\mathbb{C}}^{\hat{\mu}}$, let

$$\chi_h\colon \mathcal{M}_{\mathbb{C}}^{\hat{\mu}} \longrightarrow \mathrm{K}_0(\mathrm{HS}^{\mathrm{mon}}) \tag{3.9}$$

be the Hodge characteristic morphism defined in [4] which associates to a variety Z with a good μ_d-action the Hodge structure

$$\chi_h([Z]) = \sum_{j\in\mathbb{Z}} (-1)^j [H_c^j(Z;\mathbb{Q})] \in \mathrm{K}_0(\mathrm{HS}^{\mathrm{mon}}) \tag{3.10}$$

with the actions induced by the one $z \longmapsto \exp(2\pi\sqrt{-1}/d)z$ $(z \in Z)$ on Z. Then as in [18, Theorem 4.4], by applying the proof of [3, Theorem 4.2.1] to our situation (3.2) and (3.4), we obtain the following result.

Theorem 3.3. *In the Grothendieck group* $\mathrm{K}_0(\mathrm{HS}^{\mathrm{mon}})$*, we have*

$$[H_f^\infty] = \chi_h(\mathcal{S}_f^\infty). \tag{3.11}$$

By using Newton polyhedrons at infinity, we can rewrite Theorem 3.3 more explicitly as follows. Let $f \in \mathbb{C}[x_1, \dots, x_n]$ be a "non-convenient" polynomial such that $\dim \Gamma_\infty(f) = n$. Assume that f is non-degenerate at infinity. Now let us consider $\mathbb{C}^n$ as a toric variety associated with the fan Σ_0 in $\mathbb{R}^n$ formed by the all faces of the first quadrant $\mathbb{R}^n_+ \subset \mathbb{R}^n$. Denote by $T \simeq (\mathbb{C}^*)^n$ the open dense torus in it. Let Σ_1 be a subdivision of the dual fan of $\Gamma_\infty(f)$ which contains Σ_0 as its subfan. Then we can construct a smooth subdivision Σ of Σ_1 without subdividing the cones in Σ_0 (see e.g. [24, Lemma (2.6), Chapter II, page 99]). This implies that the toric variety X_Σ associated with Σ is a smooth compactification of $\mathbb{C}^n$. Recall that T acts on X_Σ and the T-orbits are parametrized by the cones in Σ. For a cone $\sigma \in \Sigma$ denote by $T_\sigma \simeq (\mathbb{C}^*)^{n-\dim\sigma}$ the corresponding T-orbit. We have also natural affine open subsets $\mathbb{C}^n(\sigma) \simeq \mathbb{C}^n$ of X_Σ associated to n-dimensional cones σ in Σ. Let σ be an n-dimensional cone in Σ and $\{w_1, \dots, w_n\} \subset \mathbb{Z}^n$ the set of the primitive vectors on the edges of σ. Then there exists an affine open subset $\mathbb{C}^n(\sigma)$ of X_Σ such that $\mathbb{C}^n(\sigma) \simeq \mathbb{C}^n_y$ and f has the following form on it:

$$f(y) = \sum_{v \in \mathbb{Z}^n_+} a_v y_1^{\langle w_1, v\rangle} \cdots y_n^{\langle w_n, v\rangle} = y_1^{b_1} \cdots y_n^{b_n} \times f_\sigma(y), \tag{3.12}$$

where we set $f = \sum_{v \in \mathbb{Z}^n_+} a_v x^v$,

$$b_i = \min_{v \in \Gamma_\infty(f)} \langle w_i, v\rangle \leq 0 \qquad (i = 1, 2, \dots, n) \tag{3.13}$$

and $f_\sigma(y)$ is a polynomial on $\mathbb{C}^n(\sigma) \simeq \mathbb{C}^n_y$. In $\mathbb{C}^n(\sigma) \simeq \mathbb{C}^n_y$ the hypersurface $Z := \overline{f^{-1}(0)} \subset X_\Sigma$ is explicitly written as $\{y \in \mathbb{C}^n(\sigma) \mid f_\sigma(y) = 0\}$. The variety X_Σ is covered by such affine open subsets. Let τ be a d-dimensional face of the n-dimensional cone $\sigma \in \Sigma$. For simplicity, assume that $w_1, \dots, w_d$ generate τ. Then in the affine chart $\mathbb{C}^n(\sigma) \simeq \mathbb{C}^n_y$ the T-orbit T_τ associated to τ is explicitly defined by

$$T_\tau = \{(y_1, \dots, y_n) \in \mathbb{C}^n(\sigma) \mid y_1 = \cdots = y_d = 0,\ y_{d+1}, \dots, y_n \neq 0\} \simeq (\mathbb{C}^*)^{n-d}.$$

Hence we have

$$X_\Sigma = \bigcup_{\dim\sigma = n} \mathbb{C}^n(\sigma) = \bigsqcup_{\tau \in \Sigma} T_\tau. \tag{3.14}$$

Now f was extended to a meromorphic function $\widetilde{f}$ on X_Σ, but $\widetilde{f}$ has still points of indeterminacy. From now on, we will eliminate such points by blowing up X_Σ (see [16, Section 3] and [18, Section 3] for the details). For a cone σ in Σ by taking a non-zero vector u in the relative interior rel.int(σ) of σ we define a face $\gamma(\sigma)$ of $\Gamma_\infty(f)$ by

$$\gamma(\sigma) = \left\{ v \in \Gamma_\infty(f) \mid \langle u, v\rangle = \min_{w \in \Gamma_\infty(f)} \langle u, w\rangle \right\}. \tag{3.15}$$

This face $\gamma(\sigma)$ does not depend on the choice of $u \in \mathrm{rel.int}(\sigma)$ and is called the supporting face of σ in $\Gamma_\infty(f)$. Following [14], we say that a T-orbit T_σ in X_Σ (or a cone $\sigma \in \Sigma$) is at infinity if its supporting face $\gamma(\sigma) \prec \Gamma_\infty(f)$ is at infinity i.e. $0 \notin \gamma(\sigma)$. We can easily see that $\widetilde{f}$ has poles on the union of T-orbits at infinity. Let $\rho_1, \rho_2, \dots, \rho_m$ be the 1-dimensional cones at infinity in Σ and set $T_i = T_{\rho_i}$. We call the cones ρ_i rays at infinity in Σ. Then $T_1, T_2, \dots, T_m$ are the $(n-1)$-dimensional T-orbits at infinity in X_Σ. For any $i = 1, 2, \dots, m$ the toric divisor $D_i := \overline{T_i}$ is a smooth hypersurface in X_Σ and the poles of $\widetilde{f}$ are contained in $D_1 \cup \cdots \cup D_m$. Let us denote the (unique non-zero) primitive vector in $\rho_i \cap \mathbb{Z}^n$ by u_i. Then the order $a_i > 0$ of the pole of $\widetilde{f}$ along D_i is given by

$$a_i = - \min_{v \in \Gamma_\infty(f)} \langle u_i, v \rangle. \tag{3.16}$$

Moreover by the non-degeneracy at infinity of f, the hypersurface $Z = \overline{f^{-1}(0)}$ in X_Σ intersects $D_I := \bigcap_{i \in I} D_i$ transversally for any non-empty subset $I \subset \{1, 2, \dots, m\}$. At such intersection points, $\widetilde{f}$ has indeterminacy. Now, in order to eliminate the indeterminacy of the meromorphic function $\widetilde{f}$ on X_Σ, we first consider the blow-up $\pi_1 \colon X_\Sigma^{(1)} \longrightarrow X_\Sigma$ of X_Σ along the $(n-2)$-dimensional smooth subvariety $D_1 \cap Z$. Then the indeterminacy of the pull-back $\widetilde{f} \circ \pi_1$ of $\widetilde{f}$ to $X_\Sigma^{(1)}$ is improved. If $\widetilde{f} \circ \pi_1$ still has points of indeterminacy on the intersection of the exceptional divisor E_1 of π_1 and the proper transform $Z^{(1)}$ of Z, we construct the blow-up $\pi_2 \colon X_\Sigma^{(2)} \longrightarrow X_\Sigma^{(1)}$ of $X_\Sigma^{(1)}$ along $E_1 \cap Z^{(1)}$. By repeating this procedure a_1 times, we obtain a tower of blow-ups

$$X_\Sigma^{(a_1)} \xrightarrow[\pi_{a_1}]{} \cdots\cdots \xrightarrow[\pi_2]{} X_\Sigma^{(1)} \xrightarrow[\pi_1]{} X_\Sigma. \tag{3.17}$$

Then the pull-back of $\widetilde{f}$ to $X_\Sigma^{(a_1)}$ has no indeterminacy over T_1. Next we apply this construction to the proper transforms of D_2 and Z in $X_\Sigma^{(a_1)}$. Then we obtain also a tower of blow-ups

$$X_\Sigma^{(a_1)(a_2)} \longrightarrow \cdots\cdots \longrightarrow X_\Sigma^{(a_1)(1)} \longrightarrow X_\Sigma^{(a_1)} \tag{3.18}$$

and the indeterminacy of the pull-back of $\widetilde{f}$ to $X_\Sigma^{(a_1)(a_2)}$ is eliminated over $T_1 \sqcup T_2$. By applying the same construction to (the proper transforms of) $D_3, D_4, \dots, D_m$, we finally obtain a birational morphism $\pi \colon \widetilde{X_\Sigma} \longrightarrow X_\Sigma$ such that $g := \widetilde{f} \circ \pi$ has no point of indeterminacy on the whole $\widetilde{X_\Sigma}$. Note that the smooth compactification $\widetilde{X_\Sigma}$ of $\mathbb{C}^n$ thus obtained is not a toric variety any more. By eliminating the points of indeterminacy of the meromorphic extension of f to X_Σ we have constructed the commutative diagram:

$$\begin{tikzcd} \mathbb{C}^n \arrow[r, "\iota"] \arrow[d, "f"'] & \widetilde{X_\Sigma} \arrow[d, "g"] \\ \mathbb{C} \arrow[r, "j"'] & \mathbb{P}^1. \end{tikzcd} \tag{3.19}$$

Take a local coordinate h of $\mathbb{P}^1$ in a neighborhood of $\infty \in \mathbb{P}^1$ such that $\infty = \{h = 0\}$ and set $\widetilde{g} = h \circ g$, $Y = \widetilde{g}^{-1}(0) = g^{-1}(\infty) \subset \widetilde{X_\Sigma}$ and $\Omega = \mathrm{Int}(\iota(\mathbb{C}^n) \sqcup Y)$ as before. For simplicity, let us set $\widetilde{g} = \frac{1}{f}$. Then the divisor $U = Y \cap \Omega$ in Ω contains not only the proper transforms $D'_1, \dots, D'_m$ of $D_1, \dots, D_m$ in $\widetilde{X_\Sigma}$ but also the exceptional divisors of the blow-up: $\widetilde{X_\Sigma} \longrightarrow X_\Sigma$. However as in [18], these exceptional divisors are not necessary to compute the Motivic Milnor fibers at infinity of $f\colon \mathbb{C}^n \longrightarrow \mathbb{C}$ as follows. For each non-empty subset $I \subset \{1, 2, \dots, m\}$, set $D_I = \bigcap_{i \in I} D_i$,

$$D_I^\circ = D_I \setminus \left\{ \left(\bigcup_{i \notin I} D_i \right) \cup \overline{f^{-1}(0)} \right\} \subset X_\Sigma \tag{3.20}$$

and $d_I = \gcd(a_i)_{i \in I} > 0$. Then the function $\widetilde{g} = \frac{1}{f}$ is regular on D_I° and we can decompose it as $\frac{1}{f} = \widetilde{g_1} (\widetilde{g_2})^{d_I}$ globally on a Zariski open neighborhood W of D_I° in X_Σ, where $\widetilde{g_1}$ is a unit on W and $\widetilde{g_2}\colon W \longrightarrow \mathbb{C}$ is regular. Therefore we can construct an unramified Galois covering $\widetilde{D_I^\circ}$ of D_I° with a natural μ_{d_I}-action as in (3.6). Let $[\widetilde{D_I^\circ}]$ be the element of the ring $\mathcal{M}_{\mathbb{C}}^{\hat{\mu}}$ which corresponds to $\widetilde{D_I^\circ}$. Then as in [18, Theorem 4.7] we obtain the following result.

Theorem 3.4. *Assume that* $\dim \Gamma_\infty(f) = n$ *and* f *is non-degenerate at infinity. Then we have the equality*

$$\chi_h\left(\mathcal{S}_f^\infty\right) = \sum_{I \neq \emptyset} \chi_h\left((1 - \mathbb{L})^{|I|-1} [\widetilde{D_I^\circ}] \right) \tag{3.21}$$

in the Grothendieck group $\mathrm{K}_0(\mathrm{HS}^{\mathrm{mon}})$.

For a face at infinity $\gamma \prec \Gamma_\infty(f)$ of $\Gamma_\infty(f)$, by using the lattice $M_\gamma = \mathbb{Z}^n \cap \mathbb{L}(\Delta_\gamma) \simeq \mathbb{Z}^{\dim \gamma + 1}$ in $\mathbb{L}(\Delta_\gamma)$ we set $T_{\Delta_\gamma} := \mathrm{Spec}(\mathbb{C}[M_\gamma]) \simeq (\mathbb{C}^*)^{\dim \gamma + 1}$. Moreover let $\mathbb{L}(\gamma)$ be the smallest affine linear subspace of $\mathbb{R}^n$ containing γ and for $v \in M_\gamma$ define their lattice heights $\mathrm{ht}(v, \gamma) \in \mathbb{Z}$ from $\mathbb{L}(\gamma)$ in $\mathbb{L}(\Delta_\gamma)$ so that we have $\mathrm{ht}(0, \gamma) = d_\gamma > 0$. Then to the group homomorphism $M_\gamma \longrightarrow \mathbb{C}^*$ defined by $v \longmapsto \zeta_{d_\gamma}^{\mathrm{ht}(v,\gamma)}$ we can naturally associate an element $\tau_\gamma \in T_{\Delta_\gamma}$. We define a Laurent polynomial $g_\gamma = \sum_{v \in M_\gamma} b_v x^v$ on T_{Δ_γ} by

$$b_v = \begin{cases} a_v & (v \in \gamma), \\ -1 & (v = 0), \\ 0 & (\text{otherwise}), \end{cases} \tag{3.22}$$

where $f = \sum_{v \in \mathbb{Z}_+^n} a_v x^v$. Then the Newton polytope $NP(g_\gamma)$ of g_γ is Δ_γ, $\mathrm{supp} g_\gamma \subset \{0\} \sqcup \gamma$ and the hypersurface $Z_{\Delta_\gamma}^* = \{x \in T_{\Delta_\gamma} \mid g_\gamma(x) = 0\}$ is non-degenerate (see [18, Section 4]). Since $Z_{\Delta_\gamma}^* \subset T_{\Delta_\gamma}$ is invariant by the multiplication $l_{\tau_\gamma} : T_{\Delta_\gamma} \xrightarrow{\sim} T_{\Delta_\gamma}$ by τ_γ, $Z_{\Delta_\gamma}^*$ admits an action of μ_{d_γ}. We thus obtain an element $[Z_{\Delta_\gamma}^*]$ of $\mathcal{M}_{\mathbb{C}}^{\hat{\mu}}$. For a face at infinity $\gamma \prec \Gamma_\infty(f)$ let $s_\gamma > 0$ be the dimension of the minimal coordinate subspace of $\mathbb{R}^n$ containing γ and set $m_\gamma = s_\gamma - \dim \gamma - 1 \geq 0$. Finally, for $\lambda \in \mathbb{C}$ and an element $H \in \mathrm{K}_0(\mathrm{HS}^{\mathrm{mon}})$ denote by $H_\lambda \in \mathrm{K}_0(\mathrm{HS}^{\mathrm{mon}})$ the eigenvalue λ-part of H. Then by applying the proof of [18, Theorem 5.7 (i)] to the geometric situation in Proposition 2.13, we obtain the following result.

Theorem 3.5. *Assume that* $\dim \Gamma_\infty(f) = n$ *and* f *is non-degenerate at infinity. Then for any* $\lambda \notin A_f$ *we have the equality*

$$[H_f^\infty]_\lambda = \chi_h(\mathcal{S}_f^\infty)_\lambda = \sum_\gamma \chi_h((1-\mathbb{L})^{m_\gamma} \cdot [Z^*_{\Delta_\gamma}])_\lambda \tag{3.23}$$

in $\mathrm{K}_0(\mathrm{HS}^{\mathrm{mon}})$*, where in the sum* $\sum_\gamma$ *the face* γ *of* $\Gamma_\infty(f)$ *ranges through the admissible ones at infinity.* □

4. The two-dimensional case

In this section, we consider only polynomials of two variables. We assume that $f : \mathbb{C}^2 \longrightarrow \mathbb{C}$ is "non-convenient" and satisfies the condition $\dim \Gamma_\infty(f) = 2$.

Lemma 4.1. *Assume also that* $f(x,y) \in \mathbb{C}[x,y]$ *is non-degenerate at infinity. Then the generic fiber of* $f : \mathbb{C}^2 \longrightarrow \mathbb{C}$ *is connected and hence* $H^0(f^{-1}(R); \mathbb{C}) \simeq \mathbb{C}$ *for* $R \gg 0$*.*

Proof. It is suffices to show that there is no decomposition of f of the form

$$f(x,y) = \hat{f}(\tilde{f}(x,y)) \tag{4.1}$$

by polynomials $\hat{f}(t)$ and $\tilde{f}(x,y)$ such that $\deg \hat{f}(t) \geq 2$. Assume that there exists such a decomposition $f = \hat{f} \circ \tilde{f}$ and set $m = \deg \hat{f} \geq 2$. Then we have $\Gamma_\infty(f) = m\Gamma_\infty(\tilde{f})$. Take a face at infinity $\gamma \prec \Gamma_\infty(f)$ of $\Gamma_\infty(f)$ satisfying $\dim \gamma = 1$ and let $\tilde{\gamma} \prec \Gamma_\infty(\tilde{f})$ be the corresponding one of $\Gamma_\infty(\tilde{f})$ such that $\gamma = m\tilde{\gamma}$. Denote by f_γ (resp. $\tilde{f}_{\tilde{\gamma}}$) the γ-part of f (resp. the $\tilde{\gamma}$-part of $\tilde{f}$). Then we have $f_\gamma = (\tilde{f}_{\tilde{\gamma}})^m$ for $m \geq 2$. This contradicts the non-degeneracy at infinity of f. □

In particular Lemma 4.1 and the classification of open connected Riemann surfaces implies that, for any $\lambda \neq 1$, we have the concentration:

$$H^j(f^{-1}(R); \mathbb{C})_\lambda \simeq 0 \qquad (j \neq 1) \tag{4.2}$$

for the λ-parts $H^j(f^{-1}(R); \mathbb{C})_\lambda$ of the cohomology groups of the generic fiber $f^{-1}(R)$ ($R \gg 0$) of f. In Theorem 5.2 we shall generalize this result to higher dimensions.

5. Higher-dimensional cases

In this section, we consider non-convenient polynomials $f : \mathbb{C}^n \longrightarrow \mathbb{C}$ such that $\dim \Gamma_\infty(f) = n$. For $\lambda \in \mathbb{C}$ and $j \in \mathbb{Z}$ let $H^j(f^{-1}(R); \mathbb{C})_\lambda \subset H^j(f^{-1}(R); \mathbb{C})$ be the generalized eigenspace for the eigenvalue λ of the monodromy at infinity $\Phi_j^\infty \colon H^j(f^{-1}(R); \mathbb{C}) \xrightarrow{\sim} H^j(f^{-1}(R); \mathbb{C})$ ($R \gg 0$). Denote by $\Phi_{j,\lambda}^\infty$ the restriction of Φ_j^∞ to $H^j(f^{-1}(R); \mathbb{C})_\lambda$. Assuming also that f is non-degenerate at infinity, for non-atypical eigenvalues $\lambda \notin A_f$ of f we will prove the concentration

$$H^j(f^{-1}(R); \mathbb{C})_\lambda \simeq 0 \qquad (j \neq n-1) \tag{5.1}$$

for the λ-parts $H^j(f^{-1}(R); \mathbb{C})_\lambda$ of the cohomology groups of the generic fiber $f^{-1}(R)$ ($R \gg 0$) of f. This implies that the Jordan normal forms of the λ-parts $\Phi_{j,\lambda}^\infty$ of the monodromies at infinity of f can be completely determined by $\Gamma_\infty(f)$ as in [18, Section 5]. For this purpose we need the following technical result.

Proposition 5.1. *Let* $f' \in \mathbb{C}[x_1, \dots, x_n]$ *be a polynomial such that* $\dim \Gamma_\infty(f') = n$. *Assume that* f' *is non-degenerate at infinity. Define a finite subset* $C_{f'} \subset \mathbb{C}$ *of* $\mathbb{C}$ *by* $\lambda \notin C_{f'} \iff$ *for any face at infinity* $\gamma \prec \Gamma_\infty(f')$ *contained in some proper face* $\Gamma \precneqq \Gamma_\infty(f')$ *such that* $0 \in \Gamma$ *we have* $\lambda^{d_\gamma} \neq 1$. *Let* $f'|_T : T \longrightarrow \mathbb{C}$ *be the restriction of* f' *to the open dense torus* $T = (\mathbb{C}^*)^n \subset \mathbb{C}^n$ *and for* $\lambda \in \mathbb{C}$ *and* $j \in \mathbb{Z}$ *denote the generalized eigenspace for the eigenvalue* λ *of its monodromy at infinity* $H^j((f'|_T)^{-1}(R); \mathbb{C}) \xrightarrow{\sim} H^j((f'|_T)^{-1}(R); \mathbb{C})$ $(R \gg 0)$ *by* $H^j((f'|_T)^{-1}(R); \mathbb{C})_\lambda \subset H^j((f'|_T)^{-1}(R); \mathbb{C})$. *Then for any* $\lambda \notin C_{f'}$ *we have the concentration*

$$H^j((f'|_T)^{-1}(R); \mathbb{C})_\lambda \simeq 0 \qquad (j \neq n-1) \tag{5.2}$$

for the generic fiber $(f'|_T)^{-1}(R) \subset T$ $(R \gg 0)$ *of* $f'|_T$. *Moreover if* $\lambda \notin C_{f'}$ *satisfies the condition* $H^j((f'|_T)^{-1}(R); \mathbb{C})_\lambda \neq 0$, *then there exists a face at infinity* $\gamma \prec \Gamma_\infty(f')$ *of* $\Gamma_\infty(f')$ *such that* $\lambda^{d_\gamma} = 1$.

Proof. We will prove the proposition by induction on n. If $n = 1$ the assertion is obvious. Assume that we already proved it for the lower dimensions $1, 2, \dots, n-1$. Let Σ_1' be the dual fan of $\Gamma_\infty(f')$ in $\mathbb{R}^n$ and Σ' its smooth subdivision. Then the toric variety $X_{\Sigma'}$ associated to Σ' is a smooth compactification of T. By eliminating the points of indeterminacy of the meromorphic extension of $f'|_T$ to $X_{\Sigma'}$ as in Section 3 we obtain a commutative diagram:

$$\begin{tikzcd} T \arrow[r, "\iota'"] \arrow[d, "f'|_T"'] & \widetilde{X_{\Sigma'}} \arrow[d, "g'"] \\ \mathbb{C} \arrow[r, "j"'] & \mathbb{P}^1 \end{tikzcd} \tag{5.3}$$

of holomorphic maps, where j and ι' are open embeddings. Now restricting the map $g' : \widetilde{X_{\Sigma'}} \longrightarrow \mathbb{P}^1$ to $\mathbb{C} \subset \mathbb{P}^1$ we set $X' = (g')^{-1}(\mathbb{C}) = \widetilde{X_{\Sigma'}} \setminus (g')^{-1}(\infty)$. Let $\kappa' : X' \longrightarrow \mathbb{C}$ be the restriction of g' to X'. Set $D' = X' \setminus T$ and let $i_{D'} : D' \longrightarrow X'$ and $\widetilde{\iota'} : T \longrightarrow X'$ be the inclusions. Then we obtain also a commutative diagram:

$$\begin{tikzcd} T \arrow[r, "\widetilde{\iota'}"] \arrow[d, "f'|_T"'] & X' \arrow[d, "\kappa'"] \\ \mathbb{C} \arrow[r, equal] & \mathbb{C}. \end{tikzcd} \tag{5.4}$$

Note that the normal crossing divisor D' in X' is a union of T-orbits which correspond to faces $\Gamma \prec \Gamma_\infty(f')$ such that $0 \in \Gamma$. By our induction hypothesis and the results in [14] and [16, Section 3], for $\lambda \notin C_{f'}$ the monodromies at infinity of the restrictions of κ' to these T-orbits have no λ-part. On the other hand, by applying the functor $R\kappa'_* = R\kappa'_!$ to the distinguished triangle

$$\widetilde{\iota'}_! \mathbb{C}_T \longrightarrow R\widetilde{\iota'}_* \mathbb{C}_T \longrightarrow (i_{D'})_* i_{D'}^{-1}(R\widetilde{\iota'}_* \mathbb{C}_T) \longrightarrow +1 \tag{5.5}$$

we obtain a distinguished triangle

$$R(f'|_T)_! \mathbb{C}_T \longrightarrow R(f'|_T)_* \mathbb{C}_T \longrightarrow R(\kappa'|_{D'})_* i_{D'}^{-1}(R\widetilde{\iota'}_* \mathbb{C}_T) \longrightarrow +1. \tag{5.6}$$

Then by using the above description of $\kappa'|_{D'}: D' \longrightarrow \mathbb{C}$, for $\lambda \notin C_{f'}$ we can easily show that

$$\psi_{h,\lambda}(j_! R(\kappa'|_{D'})_* i_{D'}^{-1}(R\widetilde{\iota'}_* \mathbb{C}_T)) \simeq 0 \tag{5.7}$$

and there exists an isomorphism

$$\psi_{h,\lambda}(j_! R(f'|_T)_! \mathbb{C}_T) \simeq \psi_{h,\lambda}(j_! R(f'|_T)_* \mathbb{C}_T). \tag{5.8}$$

In other words, for any $\lambda \notin C_{f'}$ and $j \in \mathbb{Z}$ we have an isomorphism

$$H_c^j((f'|_T)^{-1}(R);\mathbb{C})_\lambda \simeq H^j((f'|_T)^{-1}(R);\mathbb{C})_\lambda \qquad (R \gg 0). \tag{5.9}$$

Since the generic fiber $(f'|_T)^{-1}(R) \subset T$ $(R \gg 0)$ of $f'|_T$ is affine, the left (resp. right) hand side is zero for $j < n-1$ (resp. $j > n-1$). Hence we obtain the desired concentration

$$H^j((f'|_T)^{-1}(R);\mathbb{C})_\lambda \simeq 0 \qquad (j \neq n-1) \tag{5.10}$$

for $R \gg 0$. Now the remaining assertion follows immediately from the results in [14] and [16, Section 3]. This completes the proof. □

Theorem 5.2. *Let $f \in \mathbb{C}[x_1, \dots, x_n]$ be a non-convenient polynomial such that $\dim \Gamma_\infty(f) = n$. Assume that f is non-degenerate at infinity. Then for any non-atypical eigenvalue $\lambda \notin A_f$ of f we have the concentration*

$$H^j(f^{-1}(R);\mathbb{C})_\lambda \simeq 0 \qquad (j \neq n-1) \tag{5.11}$$

for the generic fiber $f^{-1}(R) \subset \mathbb{C}^n$ $(R \gg 0)$ of f. Moreover for such λ the relative monodromy filtration of $H^{n-1}(f^{-1}(R);\mathbb{C})_\lambda$ $(R \gg 0)$ defined by h coincides with the absolute one (up to some shift).

Proof. We will freely use the notations in Section 3. For example, we consider the commutative diagram:

$$\begin{tikzcd} \mathbb{C}^n \arrow[r, "\iota"] \arrow[d, "f"'] & \widetilde{X_\Sigma} \arrow[d, "g"] \\ \mathbb{C} \arrow[r, "j"'] & \mathbb{P}^1. \end{tikzcd} \tag{5.12}$$

By restricting the map $g : \widetilde{X_\Sigma} \longrightarrow \mathbb{P}^1$ to $\mathbb{C} \subset \mathbb{P}^1$ we set $X = g^{-1}(\mathbb{C}) = \widetilde{X_\Sigma} \setminus g^{-1}(\infty)$ and $\kappa = g|_X : X \longrightarrow \mathbb{C}$. Set $D = X \setminus \mathbb{C}^n$ and let $i_D : D \longrightarrow X$ and $\widetilde{\iota} : \mathbb{C}^n \longrightarrow X$ be the inclusions. Then we obtain also a commutative diagram:

$$\begin{tikzcd} \mathbb{C}^n \arrow[r, "\widetilde{\iota}"] \arrow[d, "f"'] & X \arrow[d, "\kappa"] \\ \mathbb{C} \arrow[r, equal] & \mathbb{C}. \end{tikzcd} \tag{5.13}$$

Note that the normal crossing divisor D in X is a union of T-orbits which correspond to atypical faces $\Gamma \prec \Gamma_\infty(f)$ of $\Gamma_\infty(f)$. By Proposition 5.1 and the results in [14] and [16, Section 3], for $\lambda \notin A_f$ the monodromies at infinity of the restrictions of κ to these

T-orbits have no λ-part. On the other hand, by applying the functor $R\kappa_* = R\kappa_!$ to the distinguished triangle

$$\widetilde{\iota}_!\mathbb{C}_{\mathbb{C}^n} \longrightarrow R\widetilde{\iota}_*\mathbb{C}_{\mathbb{C}^n} \longrightarrow (i_D)_*i_D^{-1}(R\widetilde{\iota}_*\mathbb{C}_{\mathbb{C}^n}) \longrightarrow +1 \tag{5.14}$$

we obtain a distinguished triangle

$$Rf_!\mathbb{C}_{\mathbb{C}^n} \longrightarrow Rf_*\mathbb{C}_{\mathbb{C}^n} \longrightarrow R(\kappa|_D)_*i_D^{-1}(R\widetilde{\iota}_*\mathbb{C}_{\mathbb{C}^n}) \longrightarrow +1. \tag{5.15}$$

Then by using the above description of $\kappa|_D : D \longrightarrow \mathbb{C}$, for $\lambda \notin A_f$ we can easily show that

$$\psi_{h,\lambda}(j_!R(\kappa|_D)_*i_D^{-1}(R\widetilde{\iota}_*\mathbb{C}_{\mathbb{C}^n})) \simeq 0 \tag{5.16}$$

and there exists an isomorphism

$$\psi_{h,\lambda}(j_!Rf_!\mathbb{C}_{\mathbb{C}^n}) \simeq \psi_{h,\lambda}(j_!Rf_*\mathbb{C}_{\mathbb{C}^n}). \tag{5.17}$$

In other words, for any $\lambda \notin A_f$ and $j \in \mathbb{Z}$ we have an isomorphism

$$H_c^j(f^{-1}(R);\mathbb{C})_\lambda \simeq H^j(f^{-1}(R);\mathbb{C})_\lambda \qquad (R \gg 0). \tag{5.18}$$

Since the generic fiber $f^{-1}(R) \subset \mathbb{C}^n$ $(R \gg 0)$ of f is affine, the left (resp. right) hand side is zero for $j < n-1$ (resp. $j > n-1$). Hence we obtain the desired concentration

$$H^j(f^{-1}(R);\mathbb{C})_\lambda \simeq 0 \qquad (j \neq n-1) \tag{5.19}$$

for $R \gg 0$. Moreover for such λ, by (5.17) and the proof of Sabbah [27, Theorem 13.1] the relative monodromy filtration of $H^{n-1}(f^{-1}(R);\mathbb{C})_\lambda$ $(R \gg 0)$ by h is equal to the absolute one (up to some shift). This completes the proof. □

Remark 5.3. *As is clear from the proof above, Theorem 5.2 can be easily generalized to arbitrary polynomial maps $f : U \longrightarrow \mathbb{C}$ of affine algebraic varieties U. We leave the precise formulation to the reader.*

For an element $[V] \in \mathrm{K}_0(\mathrm{HS}^{\mathrm{mon}})$, $V \in \mathrm{HS}^{\mathrm{mon}}$ with a quasi-unipotent endomorphism $\Theta\colon V \xrightarrow{\sim} V$, $p, q \geq 0$ and $\lambda \in \mathbb{C}$ denote by $e^{p,q}([V])_\lambda$ the dimension of the λ-eigenspace of the morphism $V^{p,q} \xrightarrow{\sim} V^{p,q}$ induced by Θ on the (p,q)-part $V^{p,q}$ of V. Then by Theorem 5.2 we immediately obtain the following result.

Corollary 5.4. *Assume that $\dim \Gamma_\infty(f) = n$ and f is non-degenerate at infinity. Let $\lambda \notin A_f$. Then we have $e^{p,q}([H_f^\infty])_\lambda = 0$ for $(p,q) \notin [0,n-1] \times [0,n-1]$. Moreover for any $(p,q) \in [0,n-1] \times [0,n-1]$ we have the Hodge symmetry*

$$e^{p,q}([H_f^\infty])_\lambda = e^{n-1-q,n-1-p}([H_f^\infty])_\lambda. \tag{5.20}$$

By using the notations in Section 3 we obtain the following main theorem (see [18, Theorem 5.7 (ii)]).

Theorem 5.5. *Assume that $\dim \Gamma_\infty(f) = n$ and f is non-degenerate at infinity. Let $\lambda \notin A_f$ and $k \geq 1$. Then the number of the Jordan blocks for the eigenvalue λ with sizes $\geq k$ in $\Phi_{n-1}^\infty\colon H^{n-1}(f^{-1}(R);\mathbb{C}) \xrightarrow{\sim} H^{n-1}(f^{-1}(R);\mathbb{C})$ $(R \gg 0)$ is equal to*

$$(-1)^{n-1} \sum_{p+q=n-2+k,n-1+k} \left\{ \sum_{\gamma} e^{p,q}(\chi_h((1-\mathbb{L})^{m_\gamma} \cdot [Z_{\Delta_\gamma}^*]))_\lambda \right\}, \tag{5.21}$$

where in the sum $\sum_{\gamma}$ *the face* γ *of* $\Gamma_{\infty}(f)$ *ranges through the admissible ones at infinity.*

By this theorem and the results in [18, Section 2] we immediately obtain the analogues of [18, Theorems 5.9, 5.14 and 5.16]. Here we introduce only that of [18, Theorem 5.9]. Denote by $\mathrm{Cone}_{\infty}(f)$ the closed cone $\mathbb{R}_{+}\Gamma_{\infty}(f) \subset \mathbb{R}^n_{+}$ generated by $\Gamma_{\infty}(f)$. Let $q_1, \dots, q_l$ (resp. $\gamma_1, \dots, \gamma_{l'}$) be the 0-dimensional (resp. 1-dimensional) faces at infinity of $\Gamma_{\infty}(f)$ such that $q_i \in \mathrm{Int}(\mathrm{Cone}_{\infty}(f))$ (resp. the relative interior $\mathrm{rel.int}(\gamma_i)$ of γ_i is contained in $\mathrm{Int}(\mathrm{Cone}_{\infty}(f))$). For each q_i (resp. γ_i), denote by $d_i > 0$ (resp. $e_i > 0$) the lattice distance $\mathrm{dist}(q_i, 0)$ (resp. $\mathrm{dist}(\gamma_i, 0)$) of it from the origin $0 \in \mathbb{R}^n$. For $1 \leq i \leq l'$, let Δ_i be the convex hull of $\{0\} \sqcup \gamma_i$ in $\mathbb{R}^n$. Then for $\lambda \neq 1$ and $1 \leq i \leq l'$ such that $\lambda^{e_i} = 1$ we set

$$n(\lambda)_i = \sharp\{v \in \mathbb{Z}^n \cap \mathrm{rel.int}(\Delta_i) \mid \mathrm{ht}(v, \gamma_i) = k\} + \sharp\{v \in \mathbb{Z}^n \cap \mathrm{rel.int}(\Delta_i) \mid \mathrm{ht}(v, \gamma_i) = e_i - k\}, \tag{5.22}$$

where k is the minimal positive integer satisfying $\lambda = \zeta_{e_i}^k$ and for $v \in \mathbb{Z}^n \cap \mathrm{rel.int}(\Delta_i)$ we denote by $\mathrm{ht}(v, \gamma_i)$ the lattice height of v from the base γ_i of Δ_i. Then we have the following analogue of [18, Theorem 5.9].

Theorem 5.6. *Assume that* $\dim \Gamma_{\infty}(f) = n$ *and* f *is non-degenerate at infinity. Let* $\lambda \notin A_f$. *Then we have*

1. *The number of the Jordan blocks for the eigenvalue* λ *with the maximal possible size* n *in* $\Phi^{\infty}_{n-1} \colon H^{n-1}(f^{-1}(R); \mathbb{C}) \xrightarrow{\sim} H^{n-1}(f^{-1}(R); \mathbb{C})$ $(R \gg 0)$ *is equal to* $\sharp\{q_i \mid \lambda^{d_i} = 1\}$.
2. *The number of the Jordan blocks for the eigenvalue* λ *with the second maximal possible size* $n-1$ *in* Φ^{∞}_{n-1} *is equal to* $\sum_{i \colon \lambda^{e_i}=1} n(\lambda)_i$.

Moreover in the situation above, we can obtain also a closed formula for the multiplicities of the non-atypical eigenvalues $\lambda \notin A_f$ in the monodromy at infinity $\Phi^{\infty}_{n-1} \colon H^{n-1}(f^{-1}(R); \mathbb{C}) \xrightarrow{\sim} H^{n-1}(f^{-1}(R); \mathbb{C})$ $(R \gg 0)$ as follows. We define the monodromy zeta function at infinity $\zeta^{\infty}_f(t) \in \mathbb{C}((t))$ of f by

$$\zeta^{\infty}_f(t) = \prod_{j=0}^{n-1} \det(\mathrm{id} - t\Phi^{\infty}_j)^{(-1)^j} \in \mathbb{C}((t)). \tag{5.23}$$

Then by our new compactification $\widetilde{X_{\Sigma}}$ of $\mathbb{C}^n$ we obtain the following refinement of the previous results in [14] and [16]. In particular here we can remove the condition $(*)$ in [16]. For a face at infinity $\gamma \prec \Gamma_{\infty}(f)$ of $\Gamma_{\infty}(f)$ let $\mathbb{L}(\gamma) \simeq \mathbb{R}^{\dim \gamma}$ be the minimal affine subspace of $\mathbb{R}^n$ containing γ.

Theorem 5.7. *Assume that* $\dim \Gamma_{\infty}(f) = n$ *and* f *is non-degenerate at infinity. Then we have*

$$\zeta^{\infty}_f(t) = \prod_{\gamma} (1 - t^{d_{\gamma}})^{(-1)^{s_{\gamma}-1}\mathrm{Vol}_{\mathbb{Z}}(\gamma)} \in \mathbb{C}((t)), \tag{5.24}$$

where in the product $\prod_{\gamma}$ *the face* $\gamma \prec \Gamma_{\infty}(f)$ *of* $\Gamma_{\infty}(f)$ *ranges through those at infinity satisfying the condition* $m_{\gamma} = s_{\gamma} - \dim \gamma - 1 = 0$ *and* $\mathrm{Vol}_{\mathbb{Z}}(\gamma) \in \mathbb{Z}_{>0}$ *is the normalized* $(\dim \gamma)$*-dimensional volume of* γ *with respect to the lattice* $\mathbb{L}(\gamma) \cap \mathbb{Z}^n \simeq \mathbb{Z}^{\dim \gamma}$.

Example 5.8. *Let $n = 3$ and consider a non-convenient polynomial $f(x,y,z)$ on $\mathbb{C}^3$ whose Newton polyhedron at infinity $\Gamma_\infty(f)$ is the convex hull of the points $(2,0,0),(0,2,0),(1,1,1) \in \mathbb{R}^3_+$ and the origin $0 = (0,0,0) \in \mathbb{R}^3$. Then the line segment connecting the point $(2,0,0)$ and the origin $0 \in \mathbb{R}^3$ is an atypical face of $\Gamma_\infty(f)$. Hence the 0-dimensional face at infinity $\gamma = \{(2,0,0)\} \prec \Gamma_\infty(f)$ of $\Gamma_\infty(f)$ contained in it is not admissible. However it satisfies the condition $m_\gamma = s_\gamma - \dim\gamma - 1 = 1-0-1 = 0$. For the proof of Theorem 5.7 we have to consider also the contribution from such non-admissible faces at infinity of $\Gamma_\infty(f)$.*

If we restrict ourselves to the non-atypical eigenvalues $\lambda \notin A_f$ for which we have the concentration

$$H^j(f^{-1}(R);\mathbb{C})_\lambda \simeq 0 \qquad (j \neq n-1) \tag{5.25}$$

$(R \gg 0)$ in Theorem 5.2, we have the following result.

Corollary 5.9. *Assume that $\dim\Gamma_\infty(f) = n$ and f is non-degenerate at infinity. Then for any $\lambda \notin A_f$ the multiplicity of the eigenvalue λ in the monodromy at infinity Φ^∞_{n-1} of f is equal to that of the factor $(1-\lambda t) = \lambda\cdot(1/\lambda - t)$ in the rational function*

$$\prod_\gamma (1-t^{d_\gamma})^{(-1)^{n-s_\gamma}\mathrm{Vol}_\mathbb{Z}(\gamma)} \in \mathbb{C}((t)), \tag{5.26}$$

where in the product $\prod_\gamma$ the face $\gamma \prec \Gamma_\infty(f)$ of $\Gamma_\infty(f)$ ranges through the admissible ones at infinity satisfying the condition $m_\gamma = s_\gamma - \dim\gamma - 1 = 0$. □

6. Monodromies around singular fibers

Let $f: U \longrightarrow \mathbb{C}$ be a polynomial map of an affine algebraic variety U and $B_f \subset \mathbb{C}$ the set of its bifurcation points. For a point $b \in B_f$ we choose sufficiently small $\varepsilon > 0$ such that

$$B_f \cap \{x \in \mathbb{C} \mid |x-b| \leq \varepsilon\} = \{b\} \tag{6.1}$$

and set $C_\varepsilon(b) = \{x \in \mathbb{C} \mid |x-b| = \varepsilon\} \subset \mathbb{C}$. Then we obtain a locally trivial fibration $f^{-1}(C_\varepsilon(b)) \longrightarrow C_\varepsilon(b)$ over the small circle $C_\varepsilon(b) \subset \mathbb{C}$ and the monodromy automorphisms

$$\Phi^b_j\colon H^j(f^{-1}(b+\varepsilon);\mathbb{C}) \xrightarrow{\sim} H^j(f^{-1}(b+\varepsilon);\mathbb{C}) \ \ (j = 0,1,\ldots) \tag{6.2}$$

around the central fiber $f^{-1}(b) \subset U$ associated to it. We can construct Φ^b_j's functorially as follows. Let h^b be a holomorphic local coordinate of $\mathbb{C}$ on a neighborhood of $b \in B_f$ such that $b = \{h^b(x) = 0\}$. Then to the object $\psi_{h^b}(Rf_!\mathbb{C}_U) \in \mathbf{D}^b_c(\{b\})$ and the semisimple part of the monodromy automorphism acting on it, we can associate an element

$$[H^b_f] \in \mathrm{K}_0(\mathrm{HS}^{\mathrm{mon}}). \tag{6.3}$$

Recall that the weight filtration of $[H^b_f]$ is a relative one. In this situation, we can apply our methods in previous sections to the Jordan normal forms of Φ^b_j. For the sake of simplicity, let us assume here that the central fiber $f^{-1}(b) \subset U$ is reduced and has only isolated singular points $p_1, p_2, \ldots, p_l \in f^{-1}(b)$. When f is not tame at infinity, we have to consider also the singularities at infinity of f. For this purpose, let X be a smooth compactification of U for which there exists a commutative diagram

$$\begin{tikzcd} U \arrow[r, "\iota"] \arrow[d, "f"'] & X \arrow[d, "g"] \\ \mathbb{C} \arrow[r, "j"'] & \mathbb{P}^1 \end{tikzcd} \tag{6.4}$$

of holomorphic maps. Here ι and j are are inclusion maps and g is proper. We may assume also that the divisor at infinity $D = X \setminus U \subset X$ is normal crossing and all its irreducible components are smooth. We call the irreducible components of D contained in $g^{-1}(\infty) \subset D$ (resp. in $\overline{D \setminus g^{-1}(\infty)} \subset D$) vertical (resp. horizontal) divisors at infinity of f in X. For the normal crossing divisor D let us consider the standard (minimal) stratification. Then for simplicity we assume also that the restriction $g|_{D \setminus g^{-1}(\infty)} : D \setminus g^{-1}(\infty) \longrightarrow \mathbb{C}$ of g to the horizontal part $D \setminus g^{-1}(\infty)$ of D has only stratified isolated singular points $p_{l+1}, \dots, p_{l+r}$ in $g^{-1}(b) \subset X$ and all of them are contained in the smooth part of $D \setminus g^{-1}(\infty)$. By our assumption on $f^{-1}(b) \subset U$ this implies that the hypersurface $\overline{f^{-1}(b)} = g^{-1}(b) \subset X$ in X has also an isolated singular point at each point p_i $(l+1 \leq i \leq l+r)$.

Remark 6.1. *If $U = \mathbb{C}^n$, $b \neq f(0)$ and in addition to the conditions in Theorems 5.2 and 5.5 (i.e. $\dim \Gamma_\infty(f) = n$ and f is non-degenerate at infinity) we assume that for any atypical face $\gamma \prec \Gamma_\infty(f)$ such that $\dim \gamma < n-1$ the γ-part $f_\gamma : (\mathbb{C}^*)^n \longrightarrow \mathbb{C}$ of f does not have the critical value b, then the meromorphic extension g of f to the compactification $X = \widetilde{X_\Sigma}$ satisfies the above-mentioned property in general (see also Némethi-Zaharia* [22] *and Zaharia* [35] *etc.). In this case the stratified isolated singular points $p_{l+1}, \dots, p_{l+r}$ are on the $(n-1)$-dimensional T-orbits which correspond to the atypical facets of $\Gamma_\infty(f)$.*

For $l+1 \leq i \leq l+r$, in a neighborhood of p_i the divisor D is smooth and the function $g : D \simeq \mathbb{C}^{n-1} \longrightarrow \mathbb{C}$ has an isolated singular point at $p_i \in D$. Therefore we may consider the (local) Milnor monodromies of $g : D \simeq \mathbb{C}^{n-1} \longrightarrow \mathbb{C}$ at $p_i \in D$. Denote by $A_{f,b} \subset \mathbb{C}$ the union of their eigenvalues and $1 \in \mathbb{C}$. Then by applying the proof of Theorem 5.2 to this situation, we obtain the following result. For $\lambda \in \mathbb{C}$ and $j \in \mathbb{Z}$ let $H^j(f^{-1}(b+\varepsilon); \mathbb{C})_\lambda \subset H^j(f^{-1}(b+\varepsilon); \mathbb{C})$ be the generalized eigenspace for the eigenvalue λ of the monodromy Φ_j^b around $f^{-1}(b)$.

Theorem 6.2. *In the situation as above, for any $\lambda \notin A_{f,b}$ we have the concentration*

$$H^j(f^{-1}(b+\varepsilon); \mathbb{C})_\lambda \simeq 0 \qquad (j \neq n-1). \tag{6.5}$$

Moreover for such λ the relative monodromy filtration of $H^{n-1}(f^{-1}(b+\varepsilon); \mathbb{C})_\lambda$ defined by h^b coincides with the absolute one (up to some shift). □

Corollary 6.3. *Let $\lambda \notin A_{f,b}$. Then we have $e^{p,q}([H_f^b])_\lambda = 0$ for $(p,q) \notin [0, n-1] \times [0, n-1]$. Moreover for any $(p,q) \in [0, n-1] \times [0, n-1]$ we have the Hodge symmetry*

$$e^{p,q}([H_f^b])_\lambda = e^{n-1-q, n-1-p}([H_f^b])_\lambda. \tag{6.6}$$

□

From now on, we shall use Theorem 6.2 and Corollary 6.3 to describe explicitly the Jordan normal form of Φ_{n-1}^b in terms of some Newton polyhedra associated to f. For this purpose, assume moreover that for any $1 \leq i \leq l+r$ there exists a local coordinate $y = (y_1, y_2, \dots, y_n)$ of X on a neighborhood W_i of p_i such that $p_i = \{y = 0\}$ and the local defining polynomial $f_i(y) \in \mathbb{C}[y_1, \dots, y_n]$ of the hypersurface $\overline{f^{-1}(b)} = g^{-1}(b)$ (for which we have $\overline{f^{-1}(b)} = \{f_i(y) = 0\}$) is convenient and non-degenerate at $y = 0$ (see [33] etc.). We assume also that for $l+1 \leq i \leq l+r$ we have $D = \{y_n = 0\}$ in W_i. For $1 \leq i \leq l+r$ let $\Gamma_+(f_i) \subset \mathbb{R}_+^n$ be the Newton polyhedron of f_i at $y = 0$. Moreover for $l+1 \leq i \leq l+r$ we set

$$\Gamma_+^\circ(f_i) = \Gamma_+(f_i) \cap \{v = (v_1, \dots, v_n) \in \mathbb{R}^n \mid v_n = 0\}. \tag{6.7}$$

Note that $\Gamma_+^\circ(f_i)$ is nothing but the Newton polyhedron of the restriction $f_i|_D$ of f_i to $D = \{y_n = 0\}$.

Definition 6.4. *In the situation as above, we say that a complex number $\lambda \in \mathbb{C}$ is an atypical eigenvalue for Φ_{n-1}^b if there exists a compact face $\gamma \prec \Gamma_+^\circ(f_i)$ of $\Gamma_+^\circ(f_i)$ for some $l+1 \leq i \leq l+r$ such that $\lambda^{d_\gamma} = 1$. We denote by $A_{f,b}^\circ \subset \mathbb{C}$ the set of the atypical eigenvalues for Φ_{n-1}^b.*

By the main theorem of Varchenko [33] we have $A_{f,b} \subset A_{f,b}^\circ$. On the other hand, as in [4], [18] and [19], for $1 \leq i \leq l+r$ by a toric modification $\pi_i : Y_i \longrightarrow W_i$ of W_i we can explicitly construct the motivic Milnor fiber $\mathcal{S}_{f_i,p_i} \in \mathcal{M}_{\mathbb{C}}^{\hat{\mu}}$ of f_i at p_i. See [19] for the details. For $l+1 \leq i \leq l+r$ let $(W_i \cap D)' \subset Y_i$ be the proper transform of $W_i \cap D = \{y_n = 0\}$ by π_i and $\mathcal{S}_{f_i,p_i}^\circ \in \mathcal{M}_{\mathbb{C}}^{\hat{\mu}}$ the base change of $\mathcal{S}_{f_i,p_i}$ by the inclusion map $Y_i \setminus (W_i \cap D)' \hookrightarrow Y_i$. Let $[Z_{f,b}] \in \mathcal{M}_{\mathbb{C}}^{\hat{\mu}}$ be the class of the variety $Z_{f,b} = f^{-1}(b) \setminus \{p_1, p_2, \dots, p_l\}$ with the trivial action of $\hat{\mu}$ and set

$$\mathcal{S}_f^b = [Z_{f,b}] + \sum_{i=1}^{l} \mathcal{S}_{f_i,p_i} + \sum_{i=l+1}^{l+r} \mathcal{S}_{f_i,p_i}^\circ \in \mathcal{M}_{\mathbb{C}}^{\hat{\mu}}. \tag{6.8}$$

Then as in [18, Theorem 4.4], by the proof of [3, Theorem 4.2.1] we obtain the following result.

Theorem 6.5. *In $\mathrm{K}_0(\mathrm{HS}^{\mathrm{mon}})$ we have the equality*

$$[H_f^b] = \chi_h(\mathcal{S}_f^b). \tag{6.9}$$

□

By Theorems 6.2 and 6.5 and Corollary 6.3, for any $\lambda \notin A_{f,b}^\circ$ we can describe explicitly the λ-part of the Jordan normal form of Φ_{n-1}^b as follows. For $1 \leq i \leq l+r$ let $\gamma \prec \Gamma_+(f_i)$ be a compact face of $\Gamma_+(f_i)$. Denote by Δ_γ the convex hull of $\{0\} \sqcup \gamma$ in $\mathbb{R}^n$. Let $\mathbb{L}(\Delta_\gamma)$ be the $(\dim \gamma + 1)$-dimensional linear subspace of $\mathbb{R}^n$ spanned by Δ_γ and consider the lattice $M_\gamma = \mathbb{Z}^n \cap \mathbb{L}(\Delta_\gamma) \simeq \mathbb{Z}^{\dim \gamma + 1}$ in it. Then we set $T_{\Delta_\gamma} := \mathrm{Spec}(\mathbb{C}[M_\gamma]) \simeq (\mathbb{C}^*)^{\dim \gamma + 1}$. Moreover for the points $v \in M_\gamma$ we define their lattice heights $\mathrm{ht}(v, \gamma) \in \mathbb{Z}$ from the affine hyperplane $\mathbb{L}(\gamma)$ in $\mathbb{L}(\Delta_\gamma)$ so that we have $\mathrm{ht}(0, \gamma) = d_\gamma > 0$. Then to the group homomorphism $M_\gamma \longrightarrow \mathbb{C}^*$ defined by $v \longmapsto \zeta_{d_\gamma}^{-\mathrm{ht}(v,\gamma)}$ we can naturally associate an element $\tau_\gamma \in T_{\Delta_\gamma}$. We define a Laurent polynomial $g_\gamma = \sum_{v \in M_\gamma} b_v x^v$ on T_{Δ_γ} by

$$b_v = \begin{cases} a_v & (v \in \gamma), \\ -1 & (v = 0), \\ 0 & (\text{otherwise}), \end{cases} \tag{6.10}$$

where $f = \sum_{v \in \mathbb{Z}_+^n} a_v x^v$. Then we have $NP(g_\gamma) = \Delta_\gamma$, $\mathrm{supp} g_\gamma \subset \{0\} \sqcup \gamma$ and the hypersurface $Z^*_{\Delta_\gamma} = \{x \in T_{\Delta_\gamma} \mid g_\gamma(x) = 0\}$ is non-degenerate by [18, Proposition 5.3]. Moreover $Z^*_{\Delta_\gamma} \subset T_{\Delta_\gamma}$ is invariant by the multiplication $l_{\tau_\gamma} : T_{\Delta_\gamma} \overset{\sim}{\longrightarrow} T_{\Delta_\gamma}$ by τ_γ, and hence we obtain an element $[Z^*_{\Delta_\gamma}]$ of $\mathcal{M}_{\mathbb{C}}^{\hat{\mu}}$. Finally we define $m_\gamma \in \mathbb{Z}_+$ as in Section 3. Then in the same way as [18, Theorem 5.7] and [19, Theorem 4.3] we obtain the following results.

Theorem 6.6. *In the situation as above, for any $\lambda \notin A^\circ_{f,b}$ we have the equality*

$$[H_f^b]_\lambda = \chi_h(\mathcal{S}_f^b)_\lambda = \sum_{i=1}^{l+r} \sum_{\gamma \prec \Gamma_+(f_i)} \chi_h((1-\mathbb{L})^{m_\gamma} \cdot [Z^*_{\Delta_\gamma}])_\lambda \tag{6.11}$$

in $\mathrm{K}_0(\mathrm{HS}^{\mathrm{mon}})$*, where in the sum $\sum_{\gamma \prec \Gamma_+(f_i)}$ for $l+1 \leq i \leq l+r$ the face γ of $\Gamma_+(f_i)$ ranges through compact ones not contained in $\Gamma_+^\circ(f_i)$.* □

Theorem 6.7. *In the situation as above, let $\lambda \notin A^\circ_{f,b}$ and $k \geq 1$. Then the number of the Jordan blocks for the eigenvalue λ with sizes $\geq k$ in Φ^b_{n-1} is equal to*

$$(-1)^{n-1} \sum_{p+q=n-2+k, n-1+k} \left\{ \sum_{i=1}^{l+r} \sum_{\gamma \prec \Gamma_+(f_i)} e^{p,q}(\chi_h((1-\mathbb{L})^{m_\gamma} \cdot [Z^*_{\Delta_\gamma}]))_\lambda \right\}, \tag{6.12}$$

where in the sum $\sum_{\gamma \prec \Gamma_+(f_i)}$ for $l+1 \leq i \leq l+r$ the face γ of $\Gamma_+(f_i)$ ranges through compact ones not contained in $\Gamma_+^\circ(f_i)$. □

By this theorem and the results in [18, Section 2], for $\lambda \notin A^\circ_{f,b}$ we immediately obtain the analogues of [18, Theorems 5.9, 5.14 and 5.16] for the λ-part of the Jordan normal form of Φ^b_{n-1}. It suffices to neglect the compact faces of $\Gamma_+^\circ(f_i)$ for $l+1 \leq i \leq l+r$. We omit the details.

REFERENCES

[1] Broughton, S. A. "Milnor numbers and the topology of polynomial hypersurfaces", *Invent. Math.*, 92 (1988): 217-241.

[2] Danilov, V. I. and Khovanskii, A. G. "Newton polyhedra and an algorithm for computing Hodge-Deligne numbers", *Math. Ussr Izvestiya*, 29 (1987): 279-298.

[3] Denef, J. and Loeser, F. "Motivic Igusa zeta functions", *J. Alg. Geom.*, 7 (1998): 505-537.

[4] Denef, J. and Loeser, F. "Geometry on arc spaces of algebraic varieties", *Progr. Math.*, 201 (2001): 327-348.

[5] Dimca, A. *Sheaves in topology*, Universitext, Springer-Verlag, Berlin, 2004.

[6] Dimca, A. and Saito, M. "Some consequences of perversity of vanishing cycles", *Annales de l'Institut Fourier*, 54 (2004): 1769-1792.

[7] Esterov, A. and Takeuchi, K. "Motivic Milnor fibers over complete intersection varieties and their virtual Betti numbers", *Int. Math. Res. Not.*, Vol. 2012, No. 15 (2012): 3567-3613.

[8] Fulton, W. *Introduction to toric varieties*, Princeton University Press, 1993.

[9] García López, R. and Némethi, A. "Hodge numbers attached to a polynomial map", *Ann. Inst. Fourier*, 49 (1999): 1547-1579.

[10] Guibert, G., Loeser, F. and Merle, M. "Iterated vanishing cycles, convolution, and a motivic analogue of a conjecture of Steenbrink", *Duke Math. J.*, 132 (2006): 409-457.

[11] Hotta, R., Takeuchi, K. and Tanisaki, T. *D-modules, perverse sheaves, and representation theory*, Birkhäuser Boston, 2008.

[12] Kashiwara, M. and Schapira, P. *Sheaves on manifolds*, Springer-Verlag, 1990.

[13] Kouchnirenko, A. G. "Polyédres de Newton et nombres de Milnor", *Invent. Math.*, 32 (1976): 1-31.

[14] Libgober, A. and Sperber, S. "On the zeta function of monodromy of a polynomial map", *Compositio Math.*, 95 (1995): 287-307.

[15] Matsui, Y. and Takeuchi, K. "Milnor fibers over singular toric varieties and nearby cycle sheaves", *Tohoku Math. J.*, 63 (2011): 113-136.

[16] Matsui, Y. and Takeuchi, K. "Monodromy zeta functions at infinity, Newton polyhedra and constructible sheaves", *Mathematische Zeitschrift*, 268 (2011): 409-439.

[17] Matsui, Y. and Takeuchi, K. "A geometric degree formula for A-discriminants and Euler obstructions of toric varieties", *Adv. in Math.*, 226 (2011): 2040-2064.

[18] Matsui, Y. and Takeuchi, K. "Monodromy at infinity of polynomial maps and Newton polyhedra, with Appendix by C. Sabbah", to appear in *Int. Math. Res. Not.*

[19] Matsui, Y. and Takeuchi, K. "Motivic Milnor fibers and Jordan normal forms of Milnor monodromies", arXiv:1202.5076v1, submitted.

[20] Matsui, Y. and Takeuchi, K. "On the sizes of the Jordan blocks of monodromies at infinity", arXiv:1202.5077v1, submitted.

[21] Milnor, J. *Singular points of complex hypersurfaces*, Princeton University Press, 1968.

[22] Némethi, A. and Zaharia A. "On the bifurcation set of a polynomial function and Newton boundary", *Publ. Res. Inst. Math. Sci.*, 26 (1990): 681-689.

[23] Oda, T. *Convex bodies and algebraic geometry. An introduction to the theory of toric varieties*, Springer-Verlag, 1988.

[24] Oka, M. *Non-degenerate complete intersection singularity*, Hermann, Paris (1997).

[25] Raibaut, M. "Fibre de Milnor motivique à l'infini", *C. R. Acad. Sci. Paris Sér. I Math.*, 348 (2010): 419-422.

[26] Sabbah, C. "Monodromy at infinity and Fourier transform", *Publ. Res. Inst. Math. Sci.*, 33 (1997): 643-685.

[27] Sabbah, C. "Hypergeometric periods for a tame polynomial", *Port. Math.*, 63 (2006): 173-226.

[28] Siersma, D. and Tibăr, M. "Singularities at infinity and their vanishing cycles", *Duke Math. J.*, 80 (1995): 771-783.

[29] Steenbrink, J. H. M. "Mixed Hodge structures on the vanishing cohomology", *Real and Complex Singularities*, Sijthoff and Noordhoff, Alphen aan den Rijn, (1977): 525-563.

[30] Steenbrink, J. H. M. and Zucker, S. "Variation of mixed Hodge structure I", *Invent. Math.*, 80 (1985): 489-542.

[31] Tibăr, M. "Topology at infinity of polynomial mappings and Thom regularity condition", *Compositio Math.*, 111, no.1 (1998), 89-109.

[32] Tibăr, M. *Polynomials and vanishing cycles*, Cambridge University Press, 2007.

[33] Varchenko, A. N. "Zeta-function of monodromy and Newton's diagram", *Invent. Math.*, 37 (1976): 253-262.

[34] Voisin, C. *Hodge theory and complex algebraic geometry, I*, Cambridge University Press, 2007.

[35] Zaharia A. "On the bifurcation set of a polynomial function and Newton boundary II", *Kodai Math. J.*, 19 (1996): 218-233.

Institute of Mathematics, University of Tsukuba, 1-1-1, Tennodai, Tsukuba, Ibaraki, 305-8571, Japan.
E-mail address: `takemicro@nifty.com`

Mathématiques, Laboratoire Paul Painlevé, Université Lille 1, 59655 Villeneuve d'Ascq, France.
E-mail address: `tibar@math.univ-lille1.fr`